

**Table 2.11C Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2012 and 2013**

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOTAL</b>	939	942	933	875	882	835
12	16	16	11	13	6	8
13	26	26	20	19	13	13
14	36	39	28	30	18	20
15	42	41	36	36	24	26
16	53	51	44	43	33	31
17	59	57	50	48	41	39
18	117	87	96	76	78	63
19	99	96	89	81	72	67
20	110	96	94	84	78	74
21	103	95	85	83	75	70
22	102	103	87	80	79	71
23	96	102	78	82	68	75
24	89	102	72	84	64	72
25	99	104	78	70	69	61
26-29	391	417	288	326	258	278
30-34	432	463	303	324	276	290
35-39	440	392	308	257	272	240
40-44	410	438	275	277	256	248
45-49	412	395	290	271	280	259
50-54	628	686	398	400	374	358
55-59	605	582	366	349	347	303
60-64	534	503	279	266	267	249
65 or Older	435	497	375	344	354	335

\*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.11D Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2012 and 2013**

<b>Age Category</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.36	0.36	0.36	0.33	0.34	0.32
12	0.37	0.42	0.27	0.34	0.16	0.20
13	0.62	0.58	0.49	0.43	0.33	0.31
14	0.79	0.79	0.63	0.62	0.42	0.44
15	0.86	0.88	0.77	0.82	0.54	0.61
16	0.98	0.97	0.90	0.88	0.73	0.69
17	1.04	1.13	0.96	0.99	0.82	0.84
18	1.43	1.28	1.27	1.23	1.17	1.17
19	1.35	1.29	1.35	1.28	1.28	1.22
20	1.31	1.18	1.34	1.25	1.31	1.23
21	1.20	1.19	1.31	1.35	1.25	1.24
22	1.17	1.20	1.23	1.28	1.29	1.24
23	1.13	1.10	1.24	1.24	1.21	1.23
24	1.10	1.23	1.23	1.38	1.23	1.32
25	1.15	1.25	1.37	1.31	1.32	1.25
26-29	1.16	1.18	1.19	1.28	1.18	1.19
30-34	1.01	1.04	1.14	1.17	1.10	1.12
35-39	1.20	1.13	1.26	1.12	1.16	1.09
40-44	1.08	1.07	1.17	1.08	1.14	1.02
45-49	1.09	1.06	1.14	1.09	1.11	1.09
50-54	1.48	1.35	1.44	1.47	1.39	1.36
55-59	1.42	1.60	1.50	1.46	1.45	1.32
60-64	1.68	1.67	1.43	1.38	1.38	1.31
65 or Older	1.04	1.14	0.90	0.79	0.85	0.77

\*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.11P Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2013 Versus 2012**

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOTAL</b>	0.9709		0.0278		0.0050	
12	0.4104		0.7776		0.9039	
13	0.1922		0.1512		0.1768	
14	0.3465		0.1276		0.5083	
15	0.1785		0.6380		0.3893	
16	0.0049		0.0156		0.0308	
17	0.2539		0.1510		0.0802	
18	0.0016		0.0072		0.0185	
19	0.7658		0.9892		0.5565	
20	0.2171		0.5907		0.9147	
21	0.8413		0.9578		0.3877	
22	0.4713		0.0245		0.0408	
23	0.5273		0.1834		0.0873	
24	0.5606		0.9871		0.1903	
25	0.3970		0.4245		0.5722	
26-29	0.8694		0.9040		0.3753	
30-34	0.0740		0.0459		0.0615	
35-39	0.2152		0.1381		0.1108	
40-44	0.6632		0.2930		0.2795	
45-49	0.1302		0.7269		0.4215	
50-54	0.0922		0.7189		0.4116	
55-59	0.2242		0.1361		0.0374	
60-64	0.2559		0.8715		0.6077	
65 or Older	0.3526		0.2536		0.5293	

\*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.12C Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2012 and 2013**

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOTAL</b>	956	966	890	830	839	798
12	14	11	10	10	6	7
13	21	22	17	16	11	10
14	30	31	23	22	15	14
15	40	35	32	31	20	21
16	49	45	39	36	29	26
17	56	51	46	41	37	33
18	108	75	86	63	70	54
19	91	87	78	74	66	62
20	100	89	82	77	69	69
21	92	88	75	74	66	60
22	96	93	79	75	69	67
23	90	98	70	75	64	66
24	85	97	66	80	60	67
25	97	97	71	64	63	58
26-29	374	405	261	299	237	260
30-34	410	439	281	293	260	263
35-39	427	375	281	235	249	222
40-44	389	411	239	250	227	230
45-49	393	371	268	238	258	232
50-54	601	661	366	350	347	338
55-59	598	555	312	292	301	279
60-64	516	480	258	232	243	224
65 or Older	454	516	341	311	320	303

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.12D Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2012 and 2013**

<b>Age Category</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.37	0.37	0.34	0.32	0.32	0.30
12	0.35	0.30	0.25	0.27	0.15	0.17
13	0.52	0.52	0.41	0.38	0.27	0.23
14	0.68	0.66	0.53	0.47	0.35	0.32
15	0.83	0.78	0.70	0.72	0.46	0.51
16	0.96	0.91	0.83	0.77	0.66	0.60
17	1.03	1.06	0.90	0.88	0.76	0.74
18	1.46	1.31	1.32	1.18	1.12	1.05
19	1.34	1.35	1.30	1.32	1.23	1.25
20	1.31	1.26	1.28	1.25	1.26	1.24
21	1.23	1.28	1.26	1.31	1.23	1.15
22	1.22	1.23	1.23	1.34	1.24	1.25
23	1.18	1.20	1.22	1.27	1.18	1.19
24	1.15	1.27	1.25	1.36	1.20	1.27
25	1.21	1.24	1.33	1.24	1.28	1.22
26-29	1.20	1.25	1.16	1.26	1.12	1.17
30-34	1.07	1.05	1.11	1.11	1.07	1.07
35-39	1.25	1.17	1.21	1.07	1.10	1.04
40-44	1.15	1.12	1.07	1.03	1.05	0.98
45-49	1.14	1.08	1.08	1.00	1.05	1.00
50-54	1.52	1.39	1.34	1.34	1.30	1.30
55-59	1.50	1.65	1.36	1.28	1.33	1.24
60-64	1.70	1.73	1.38	1.25	1.31	1.22
65 or Older	1.08	1.19	0.81	0.72	0.77	0.70

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.12P Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2013 Versus 2012**

<b>Age Category</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.7810		0.0669		0.0411	
12	0.0428		0.6306		0.6568	
13	0.2731		0.1555		0.1780	
14	0.3740		0.0645		0.1492	
15	0.0597		0.4110		0.8419	
16	0.0044		0.0032		0.0070	
17	0.2786		0.0467		0.0798	
18	0.0000		0.0001		0.0050	
19	0.7810		0.6024		0.6498	
20	0.3833		0.5235		0.8649	
21	0.6992		0.3917		0.2430	
22	0.6195		0.0719		0.1934	
23	0.5285		0.6163		0.5348	
24	0.4774		0.8236		0.2382	
25	0.1003		0.3573		0.4998	
26-29	0.6608		0.9788		0.6700	
30-34	0.0866		0.1326		0.3996	
35-39	0.1449		0.5659		0.4675	
40-44	0.8893		0.9065		0.5306	
45-49	0.0714		0.5215		0.2704	
50-54	0.0646		0.3945		0.4206	
55-59	0.4297		0.3522		0.2918	
60-64	0.3954		0.6795		0.7814	
65 or Older	0.2021		0.6525		0.9615	

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.13C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2012 and 2013**

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOTAL</b>	705	727	325	334	298	300
12	7	10	5	8	4	4
13	15	13	11	9	6	4
14	19	22	14	19	10	12
15	23	22	17	18	12	12
16	28	27	21	24	17	15
17	30	32	23	25	17	18
18	42	44	32	35	23	27
19	50	50	38	38	25	27
20	49	46	34	35	24	28
21	47	48	30	36	24	26
22	49	59	34	33	29	24
23	53	55	31	32	24	24
24	43	54	27	29	23	23
25	61	46	30	28	24	21
26-29	206	202	109	108	97	88
30-34	212	223	103	108	92	92
35-39	229	222	112	106	99	93
40-44	231	250	120	111	110	107
45-49	212	224	94	103	88	96
50-54	284	355	99	108	95	100
55-59	285	241	101	103	97	102
60-64	214	227	85	51	79	46
65 or Older	305	367	122	125	117	124

\*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.13D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2012 and 2013**

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOTAL</b>	0.27	0.28	0.13	0.13	0.11	0.11
12	0.18	0.26	0.12	0.22	0.09	0.11
13	0.37	0.31	0.27	0.21	0.15	0.10
14	0.44	0.49	0.34	0.42	0.25	0.27
15	0.52	0.51	0.39	0.43	0.29	0.29
16	0.63	0.60	0.48	0.53	0.38	0.35
17	0.66	0.73	0.52	0.58	0.37	0.42
18	0.89	0.85	0.67	0.69	0.50	0.55
19	1.06	1.01	0.83	0.85	0.57	0.62
20	0.97	0.97	0.75	0.77	0.52	0.63
21	0.98	1.02	0.67	0.79	0.54	0.60
22	0.94	1.09	0.70	0.69	0.61	0.53
23	1.04	1.06	0.69	0.70	0.55	0.54
24	0.93	1.04	0.65	0.63	0.55	0.52
25	1.27	1.00	0.70	0.66	0.58	0.51
26-29	1.03	1.01	0.61	0.59	0.55	0.49
30-34	0.91	0.94	0.50	0.52	0.45	0.44
35-39	0.99	1.04	0.54	0.55	0.49	0.48
40-44	0.98	0.98	0.58	0.51	0.54	0.49
45-49	0.90	0.96	0.44	0.50	0.41	0.47
50-54	1.11	1.35	0.42	0.47	0.41	0.43
55-59	1.21	1.07	0.49	0.50	0.47	0.49
60-64	1.18	1.15	0.50	0.29	0.46	0.26
65 or Older	0.73	0.85	0.29	0.29	0.28	0.29

\*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.



**Table 2.13P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2013 Versus 2012**

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOTAL</b>	0.8283		0.5328		0.4649	
12	0.6948		0.1545		0.6591	
13	0.2056		0.1208		0.0662	
14	0.7970		0.8833		0.8174	
15	0.8197		0.1645		0.7467	
16	0.3275		0.9590		0.4357	
17	0.7949		0.7245		0.8261	
18	0.9270		0.3051		0.0871	
19	0.2291		0.4029		0.1437	
20	0.7483		0.1929		0.3204	
21	0.3454		0.3222		0.5314	
22	0.0301		0.6041		0.2785	
23	0.9687		0.6965		0.8843	
24	0.3070		0.6669		0.8254	
25	0.0856		0.5517		0.4631	
26-29	0.2998		0.3816		0.2015	
30-34	0.9808		0.1951		0.2137	
35-39	0.0743		0.2667		0.2617	
40-44	0.5809		0.0266		0.1142	
45-49	0.0444		0.2532		0.2873	
50-54	0.0167		0.2091		0.5408	
55-59	0.0765		0.7489		0.6302	
60-64	0.6190		0.0520		0.0367	
65 or Older	0.4884		0.4706		0.7478	

\*Low precision; no estimate reported.

NOTE: Smokeless Tobacco includes chewing tobacco or snuff.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.14C Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2012 and 2013**

<b>Age Category</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	882	922	511	493	388	354
12	7	8	5	6	3	3
13	15	12	10	11	6	4
14	21	20	18	14	10	7
15	27	24	23	21	11	12
16	36	35	28	28	18	17
17	42	39	36	32	21	23
18	74	64	62	56	42	37
19	72	69	63	56	45	36
20	84	72	63	57	39	39
21	76	76	57	58	43	38
22	79	81	55	53	38	34
23	76	78	47	50	30	38
24	67	75	46	48	30	34
25	71	75	45	43	30	29
26-29	287	312	173	184	135	122
30-34	314	345	166	180	110	122
35-39	316	296	159	128	111	91
40-44	293	315	140	147	103	101
45-49	289	279	148	131	107	98
50-54	424	488	178	238	133	155
55-59	411	394	207	185	177	98
60-64	351	351	149	144	92	117
65 or Older	413	456	135	136	105	115

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.14D Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2012 and 2013**

<b>Age Category</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.34	0.35	0.20	0.19	0.15	0.14
12	0.17	0.22	0.13	0.15	0.07	0.08
13	0.37	0.29	0.24	0.25	0.15	0.10
14	0.49	0.46	0.42	0.32	0.24	0.15
15	0.60	0.56	0.53	0.49	0.25	0.29
16	0.75	0.73	0.63	0.60	0.43	0.38
17	0.91	0.87	0.80	0.74	0.47	0.54
18	1.13	1.10	1.04	1.02	0.79	0.73
19	1.30	1.21	1.18	1.09	0.92	0.80
20	1.32	1.22	1.17	1.13	0.83	0.83
21	1.22	1.25	1.11	1.14	0.88	0.84
22	1.24	1.28	1.01	1.04	0.78	0.73
23	1.29	1.23	0.98	1.00	0.66	0.82
24	1.21	1.24	1.01	0.98	0.70	0.73
25	1.35	1.32	1.00	1.01	0.70	0.71
26-29	1.22	1.25	0.91	0.92	0.76	0.66
30-34	1.12	1.21	0.77	0.82	0.53	0.58
35-39	1.23	1.23	0.77	0.65	0.55	0.47
40-44	1.12	1.13	0.66	0.64	0.50	0.46
45-49	1.08	1.08	0.66	0.61	0.49	0.48
50-54	1.48	1.55	0.74	0.98	0.56	0.67
55-59	1.50	1.57	0.94	0.87	0.82	0.47
60-64	1.61	1.59	0.86	0.78	0.54	0.64
65 or Older	0.99	1.05	0.32	0.31	0.25	0.27

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.14P Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2013 Versus 2012**

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOTAL</b>	0.8196		0.1812		0.0351	
12	0.4169		0.5629		0.6641	
13	0.3536		0.7240		0.0501	
14	0.1348		0.0034		0.0055	
15	0.0514		0.1696		0.3816	
16	0.0032		0.0136		0.1889	
17	0.3107		0.2612		0.9956	
18	0.0049		0.0430		0.1030	
19	0.8281		0.8917		0.6432	
20	0.3869		0.5653		0.9160	
21	0.7462		0.6910		0.9958	
22	0.7975		0.0653		0.0245	
23	0.2223		0.2942		0.1726	
24	0.5823		0.5719		0.4074	
25	0.5414		0.0723		0.1141	
26-29	0.5041		0.2438		0.8848	
30-34	0.2349		0.2984		0.2292	
35-39	0.2202		0.0844		0.0289	
40-44	0.8182		0.5658		0.9810	
45-49	0.4432		0.7128		0.7461	
50-54	0.0203		0.5520		0.7959	
55-59	0.0508		0.0583		0.0017	
60-64	0.8282		0.4376		0.3277	
65 or Older	0.2803		0.3710		0.5470	

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.15C Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2012 and 2013**

<b>Age Category</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	675	801	955	1,007	1,016	1,076
12	25	22	22	16	9	8
13	32	32	28	25	17	15
14	49	50	44	39	30	25
15	55	51	48	45	31	30
16	64	66	57	58	40	40
17	74	72	69	66	52	49
18	140	114	136	108	115	87
19	125	109	122	103	94	85
20	126	113	122	107	104	92
21	120	118	118	116	108	105
22	118	122	116	117	106	106
23	112	118	107	112	97	104
24	104	114	100	108	89	99
25	109	116	104	110	99	99
26-29	456	468	435	446	385	410
30-34	470	506	441	470	389	413
35-39	492	445	461	405	414	367
40-44	469	513	443	472	386	412
45-49	485	443	425	407	380	371
50-54	697	739	627	676	564	602
55-59	684	656	587	550	516	493
60-64	599	578	530	488	466	458
65 or Older	391	462	506	533	483	533

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.15D Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2012 and 2013**

<b>Age Category</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.26	0.31	0.37	0.38	0.39	0.41
12	0.60	0.57	0.52	0.43	0.23	0.21
13	0.74	0.73	0.65	0.58	0.42	0.36
14	0.98	0.98	0.92	0.82	0.67	0.56
15	1.01	1.02	0.97	0.96	0.68	0.68
16	1.07	1.12	1.06	1.04	0.86	0.84
17	1.09	1.13	1.11	1.14	0.98	0.98
18	1.22	1.17	1.30	1.14	1.44	1.22
19	1.22	1.25	1.36	1.33	1.36	1.38
20	1.04	1.08	1.18	1.21	1.41	1.41
21	0.84	0.87	0.94	0.96	1.23	1.22
22	0.76	0.77	0.88	0.96	1.23	1.20
23	0.74	0.69	1.02	0.98	1.15	1.22
24	0.69	0.82	0.93	1.02	1.19	1.18
25	0.78	0.70	1.01	0.98	1.30	1.31
26-29	0.72	0.76	0.96	1.03	1.26	1.22
30-34	0.73	0.79	0.95	1.02	1.10	1.16
35-39	0.82	0.81	1.08	1.11	1.26	1.22
40-44	0.77	0.77	1.04	1.09	1.17	1.16
45-49	0.68	0.80	1.01	1.12	1.13	1.21
50-54	0.98	0.85	1.32	1.41	1.46	1.58
55-59	1.14	1.17	1.61	1.66	1.68	1.67
60-64	1.14	1.39	1.75	1.74	1.83	1.75
65 or Older	0.93	1.06	1.21	1.23	1.15	1.23

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.15P Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2013 Versus 2012**

<b>Age Category</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.0373		0.3796		0.9167	
12	0.4878		0.1914		0.7998	
13	0.2433		0.1896		0.5894	
14	0.0357		0.0070		0.0503	
15	0.0366		0.0999		0.1242	
16	0.4921		0.1806		0.0188	
17	0.9813		0.9083		0.4035	
18	0.1101		0.0694		0.0081	
19	0.8202		1.0000		0.6803	
20	0.1436		0.4488		0.4146	
21	0.9631		0.7165		0.8718	
22	0.4115		0.1424		0.8020	
23	0.3348		0.4227		0.4845	
24	0.5299		0.9609		0.4682	
25	0.7843		0.6343		0.2353	
26-29	0.5074		0.8745		0.2350	
30-34	0.8830		0.8957		0.3925	
35-39	0.7229		0.2017		0.2327	
40-44	0.9825		0.6548		0.7068	
45-49	0.0506		0.7233		0.9545	
50-54	0.1688		0.4423		0.5292	
55-59	0.7300		0.3563		0.7855	
60-64	0.1907		0.2108		0.8473	
65 or Older	0.1790		0.1943		0.7441	

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.16C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2012 and 2013**

Age Category	Alcohol Use (2012)	Alcohol Use (2013)	Binge Alcohol Use (2012)	Binge Alcohol Use (2013)	Heavy Alcohol Use (2012)	Heavy Alcohol Use (2013)
<b>TOTAL</b>	1,016	1,076	797	804	448	438
12	9	8	6	4	1	3
13	17	15	12	11	5	3
14	30	25	18	17	4	5
15	31	30	22	21	7	8
16	40	40	30	28	14	11
17	52	49	41	38	18	19
18	115	87	81	63	30	30
19	94	85	75	71	44	33
20	104	92	89	71	48	40
21	108	105	87	81	47	42
22	106	106	84	83	48	44
23	97	104	76	78	39	43
24	89	99	71	78	33	43
25	99	99	80	72	52	34
26-29	385	410	276	309	132	156
30-34	389	413	281	306	144	190
35-39	414	367	253	254	129	130
40-44	386	412	256	263	139	133
45-49	380	371	234	239	118	119
50-54	564	602	357	360	190	166
55-59	516	493	270	272	138	134
60-64	466	458	216	217	121	118
65 or Older	483	533	266	297	136	150

\*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.



**Table 2.16D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Standard Errors of Percentages, 2012 and 2013**

Age Category	Alcohol Use (2012)	Alcohol Use (2013)	Binge Alcohol Use (2012)	Binge Alcohol Use (2013)	Heavy Alcohol Use (2012)	Heavy Alcohol Use (2013)
<b>TOTAL</b>	0.39	0.41	0.31	0.31	0.17	0.17
12	0.23	0.21	0.14	0.11	0.02	0.07
13	0.42	0.36	0.29	0.25	0.13	0.07
14	0.67	0.56	0.41	0.39	0.09	0.10
15	0.68	0.68	0.49	0.50	0.18	0.20
16	0.86	0.84	0.66	0.62	0.34	0.25
17	0.98	0.98	0.81	0.83	0.40	0.44
18	1.44	1.22	1.17	1.05	0.59	0.57
19	1.36	1.38	1.25	1.28	0.89	0.71
20	1.41	1.41	1.46	1.28	0.96	0.85
21	1.23	1.22	1.37	1.31	0.97	0.86
22	1.23	1.20	1.34	1.29	0.94	0.88
23	1.15	1.22	1.24	1.30	0.79	0.87
24	1.19	1.18	1.26	1.28	0.77	0.89
25	1.30	1.31	1.40	1.28	1.14	0.78
26-29	1.26	1.22	1.24	1.28	0.72	0.82
30-34	1.10	1.16	1.10	1.10	0.68	0.86
35-39	1.26	1.22	1.13	1.14	0.64	0.67
40-44	1.17	1.16	1.05	1.03	0.66	0.59
45-49	1.13	1.21	0.97	1.00	0.53	0.58
50-54	1.46	1.58	1.35	1.35	0.79	0.70
55-59	1.68	1.67	1.18	1.20	0.65	0.64
60-64	1.83	1.75	1.22	1.17	0.71	0.65
65 or Older	1.15	1.23	0.64	0.68	0.33	0.34

\*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.16P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2013 Versus 2012**

Age Category	Alcohol Use (2012)	Alcohol Use (2013)	Binge Alcohol Use (2012)	Binge Alcohol Use (2013)	Heavy Alcohol Use (2012)	Heavy Alcohol Use (2013)
<b>TOTAL</b>	0.9167		0.8875		0.3092	
12	0.7998		0.3371		0.2003	
13	0.5894		0.9594		0.1115	
14	0.0503		0.1053		0.8857	
15	0.1242		0.2460		0.5556	
16	0.0188		0.0299		0.1072	
17	0.4035		0.1728		0.9483	
18	0.0081		0.0164		0.4733	
19	0.6803		0.6463		0.0255	
20	0.4146		0.6471		0.2729	
21	0.8718		0.2279		0.0606	
22	0.8020		0.7012		0.5048	
23	0.4845		0.6420		0.7985	
24	0.4682		0.4325		0.6273	
25	0.2353		0.0471		0.0113	
26-29	0.2350		0.1733		0.2116	
30-34	0.3925		0.3337		0.1406	
35-39	0.2327		0.6821		0.4583	
40-44	0.7068		0.2295		0.4211	
45-49	0.9545		0.6100		0.7913	
50-54	0.5292		0.8820		0.3558	
55-59	0.7855		0.6385		0.2441	
60-64	0.8473		0.9114		0.6507	
65 or Older	0.7441		0.3316		0.7216	

\*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.